



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07B 37/00	A1	(11) International Publication Number: WO 97/43230 (43) International Publication Date: 20 November 1997 (20.11.97)
(21) International Application Number: PCT/SE97/00794 (22) International Filing Date: 14 May 1997 (14.05.97) (30) Priority Data: 9601856-9 14 May 1996 (14.05.96) SE 9603913-6 25 October 1996 (25.10.96) SE (71) Applicant (for all designated States except US): LABWELL AB [SE/SE]; Hamnplan 9, S-753 19 Uppsala (SE). (72) Inventors; and (75) Inventors/Applicants (for US only): LINDEBERG, Gunnar [SE/SE]; Murargatan 31, S-754 37 Uppsala (SE). LARHED, Mats [SE/SE]; St. Johannesgatan 82, S-752 64 Uppsala (SE). HALLBERG, Anders [SE/SE]; Björklundavägen 4 A, S-756 46 Uppsala (SE). (74) Agents: ALDENBÄCK, Ulla et al.; Dr. Ludwig Brann Patentbyrå AB, P.O. Box 1344, S-751 43 Uppsala (SE).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: METHOD FOR ORGANIC REACTIONS		
(57) Abstract		
<p>The present invention relates to a method for organic reactions. More specifically, a method for palladium, except Pd/C, catalyzed organic reactions comprising a heating step with microwave energy. One type of organic reactions concerned are coupling reactions, in which new carbon-carbon bonds are formed. Preferred reactions are the Heck, Stille and Suzuki reactions. The method provides high yields in very short reaction times.</p>		
BEST AVAILABLE COPY		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TC	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHOD FOR ORGANIC REACTIONS

FIELD OF THE INVENTION

The present invention relates to a method for transition metal catalyzed organic reactions comprising a heating step. More precisely, the invention relates to organic reactions catalyzed with palladium, except Pd/C, under microwave energy. One type of organic reactions concerned are coupling reactions, in which new carbon-carbon bonds are formed.

BACKGROUND OF THE INVENTION

Transition metal catalyzed reactions have played a role in organic synthesis for a very long time. Twenty five years ago the first examples on the use of palladium as a catalyst for coupling reactions were disclosed. Since then enormous efforts have been devoted to extend the scope of palladium-catalyzed reactions. Among these, the Heck reaction¹, the Stille reaction² and the Suzuki reaction³ probably represent the most applied and most reliable reactions considering the reactions delivering new carbon-carbon bond formation. These reactions allow presence of a wide variety of substituents attached to the reactants and merits special attention due to the simplicity of the experimental procedures. The Heck reaction, which is a vinylic substitution reaction is most frequently conducted with olefines and organo halides or triflates as reactants. Couplings of organotin reagents with organo halides or triflates are named Stille reactions. The related couplings of organoboronic acids and/or organoboronic esters with organo halides or triflates are named Suzuki reactions. These reaction are of outmost importance in organic synthesis and have all found use in Combinatorial Chemistry (CC), sometimes as key reactions in the creation of chemical libraries⁴. Combinatorial Chemistry is conducted either in solution or preferably on solid phase. Combinatorial Chemistry combined with High Throughput Screening (HTS) has revolutionised the Drug Discovery Programmes in the pharmaceutical industry during the last few years⁵.

In Combinatorial Chemistry the reaction time factor is of importance. Rapid reactions are desired. The long reaction times often required in the palladium-catalyzed coupling reactions presented above is therefore in this respect a severe limitation. While some combinations of reactants allow fast conversions, and efficient use in Combinatorial Chemistry have been demonstrated, the majority of reactant combinations require unacceptable reaction times for completion. With regard to the palladium-catalyzed Heck, Stille or Suzuki reactions, attempted enhancement of the conversion rate by increasing the reaction temperature to over 130-150°C most

often leads to collapse of the catalytic system before a full conversion is achieved. Product decomposition is frequently observed and a mixture of undesired side products are formed.

PRIOR ART

Microwave assisted transition-metal catalyzed reactions of double bonds, including hydrogenation, catalyzed by $RaNi$ or Pd/C ,⁶ and addition of chlorinated hydrocarbons to double bonds, catalyzed by CuI ⁷ have been reported.

SUMMARY OF THE INVENTION

The present invention provides a method for palladium, except Pd/C , catalyzed organic reactions comprising a heating step performed with microwave energy.

In a preferred embodiment, the reactions are coupling reactions such as the Heck, Stille and Suzuki reactions.

Furthermore, the present invention provides for use of palladium-catalyzed coupling reactions in combinatorial chemistry and for creation of chemical libraries.

DETAILED DESCRIPTION OF THE INVENTION

1. PALLADIUM-CATALYZED COUPLINGS IN SOLUTION

Microwave assisted methodology⁸ is based on a fast heating of the reaction mixtures to high temperatures. The inventors surprisingly found that the Heck, Stille and Suzuki reactions in which labile organotransition metal intermediates play a pivotal role in the catalytic cycle, was compatible with the extreme temperatures used in these reactions. It was found that the Heck, Stille and Suzuki reactions, after very short reaction times (2-7 minutes) when applying microwave assisted technology, allowed isolation of reaction products in high yields and also with high purity, see TABLE 1 below. The corresponding reactions performed with standard heating required hours to reach satisfactory yields according to literature, as denoted in TABLE 1. While an increase of the reaction temperature in the latter cases often provided complex reaction mixtures, the microwave assisted reactions were leading to essentially no undesired side products. Preferably the microwave energy is solely or predominantly in the form of a standing wave.

The inventors have reacted aryl iodides, aryl bromides and aryl triflates with various olefins (Heck reactions). These olefins were chosen to permit comparison of microwave-assisted reactions with conventionally heated reactions, with respect to regio- and stereoselectivity, as well as double bond isomerization. The reactions were conducted in sealed pyrex vessels and a commercially available reactor producing continuous irradiation was used. Our initial screening experiments, with different solvents suggested that DMF was suitable for all reactant combinations tested. Acetonitrile, ethanol, water, DME and NMP are examples of other suitable solvents. With the exception of reaction scale, solvent and heating procedure the reactions were performed under reaction conditions identical to the original procedures as described in the references cited, in order to enable an accurate comparison. The inventors intended to utilize microwave assisted reactions in Combinatorial Chemistry and the primary objective was minimization of the reaction times. Irradiation effects were altered to allow full conversions of the arylating agents in less than seven minutes.

Methyl acrylate was smoothly converted in 3.5 min at 60 W, to the corresponding cinnamic acid ester in a standard Heck reaction medium. The same transformation could be conducted at a lower power although a longer reaction time was required. In order to achieve a full consumption of 1-iodonaphthalene in a reaction with acrylonitrile in the same reaction time, a power of 80 W was used. While methyl acrylate provided the E isomer exclusively, the reaction with acrylonitrile produced a mixture of the E and Z isomers (E/Z = 4/1) as expected from related standard reactions with this olefin. Arylation of styrene with iodobenzene delivered a terminal/internal ratio of 19/1, in accordance with the original procedure. Microwave assisted arylation of dihydrofuran according to Larock's original procedure furnished a mixture of double bond isomers. 2-Phenyl-2, 3-dihydrofuran was isolated in fair yield. A reaction time of 6 min at 30 W was needed in this case, to be compared with 24 hours at 80° C when standard heating was employed. For studies of the regioselectivity with regard to the direction of insertion, alkyl vinyl ethers were used as substrates. Procedures have been developed that allow either electronically controlled internal arylation by application of bidentate ligands or alternatively, chelation-controlled terminal arylation. Microwave assisted arylation of butyl vinyl ether with aryl triflate provided a mixture of α -arylated vinyl ether and the corresponding acetophenone. 4-Acetyl tert. butylbenzene could be isolated in good

yield after conducting the reaction in presence of a minor amount of water. The reactions of dimethylaminoethyl vinyl ether with 2-naphthyl triflate as the substrate constitutes an example of a sluggish reaction that requires 9 hours at 60° C to afford fair yields of coupled product. In the microwave assisted reaction, it was essential to employ a low effect, 35W, to avoid decomposition of the catalytic system. A longer reaction time, 7 min, was therefore needed. A highly regioselective arylation at the terminal carbon occurred, and equal amount of the E and Z isomers were formed, in full agreement with the reference procedures. Allyldimethylamine and allyltrimethylsilane were both successfully arylated at the internal olefinic carbon with high regioselectivity after 5 minutes.

Suzuki coupling of phenyl boronic acid with 4-bromotoluene as well as Stille coupling of 4-acetyl phenyl triflate occurred smoothly with microwave irradiation.

Arylation of styrene with 4-bromo-1-iodobenzene delivered a terminal/internal ratio of 15/1 in accordance with the original procedure. The result from this reaction shows that the microwave irradiation provides the same selectivity, with respect to halide displacement as compared to the original reaction and substitution of the iodoatom occurs exclusively. No products derived from bromo substitution were formed, demonstrating the potential of microwave irradiation technology for combinatorial chemistry. The bromo atom remains intact and provides a handle for further consecutive palladium-catalyzed transformations.

2. PALLADIUM-CATALYZED COUPLINGS ON SOLID SUPPORT

The inventors have found that palladium-catalyzed reactions on polymer-support were compatible with the microwave assisted methodology. Preferably the microwave energy is solely or predominantly in the form of a standing wave. High yields of products with high purity were isolated after short reaction times. No or very limited decomposition of the solid phase was observed. Examples of microwave-assisted Suzuki couplings on solid phase are demonstrated in TABLE 2 below. We coupled 4-iodo and 4-bromo- benzoic acid to Rink resin¹⁸(Tentagel¹⁹) and conducted the palladium-catalyzed reactions with aryl and heteroaryl boronic acids substituted with a variety of substituents. Both arylbromo and aryliodo groups on the resin are good coupling partners with organoboronic acids.

One example of a Stille coupling on solid phase and one example of a Heck

coupling on solid phase are shown in TABLE 3.

Entry 15 in TABLE 2 demonstrates that the microwave technology permit selective coupling of 4-bromophenylboronic acid to aryl iodide attached to the resin. The organoiodide is more reactive and the bromo atom is not displaced. A high chemoselectivity is achieved and the bromo atom constitutes a handle for consecutive reactions.

Table 1. Palladium-Catalyzed Coupling Reactions in Solution under Microwave Irradiation.















































entry	aryl halide or aryl triflate	olefine or organometallic	time effect	product	isolated yield ^a	conventional heating time (min)	yield (%)
1			3.50 min. 60 W Pd(OAc) ₂ , DMF		70%	300	68% ⁹
2			3.50 min. 60 W Pd(OAc) ₂ , DMF o-(tol) ₃ P		94%	120	70% ¹⁰
3			2.50 min. 90 W Pd(OAc) ₂ , DMF		87%	120	75% ⁹
4			3.50 min. 80 W Pd(OAc) ₂ , DMF		90%		
5			6.00 min. 30 W Pd(OAc) ₂ , Ph ₃ P DMF, Bu ₄ NC ⁺		58%	1440	76% ¹¹
6			2.50 min. 55 W Pd(OAc) ₂ , DPPP DMF, H ₂ O		77%		
7			7.00 min. 35 W Pd(OAc) ₂ , Ph ₃ P DMF		87%	540	93% ¹²

Table 1. Palladium-Catalyzed Coupling Reactions in Solution under Microwave Irradiation.


entry	aryl halide or aryl triflate	olefine or organometallic	time effect	product	isolated yield ^a	conventional heating time (min)	yield (%)
8			5.00 min. 50 W Pd(OAc) ₂ , DPPF CH ₃ CN		54%	960	59%
9			5.00 min. 50 W Pd(OAc) ₂ , DPPF CH ₃ CN		64%	2880	72%
10			2.50 min. 55 W Pd(PPh ₃) ₄ , EtOH, DME, H ₂ O		55%	360	94% ¹³
11			2.50 min. 50 W Pd ₂ (dba) ₃ , Ph ₃ As, LiCl, NMP		68%	8400	82% ¹⁴
12			4.50 min. 60 W Pd(OAc) ₂ , DMF		63%	1020	64% ¹⁵

a) Purity >95% by GC/MS

Table 2. Suzuki Coupling Reactions on Solid Phase under Microwave Irradiation^a.

			
Entry		Entry	
1, 2	 98% ; X=I 98% ; X=Br	10	 89% ; X=I
3, 4	 96% ; X=I 95% ; X=Br	11, 12	 86% ; X=I 84% ; X=Br
5	 88% ; X=I	13, 14	 89% ; X=I 88% ; X=Br
6, 7	 99% ; X=I 93% ; X=Br	15 ^b	 90% ; X=I
8, 9	 99% ; X=I 96% ; X=Br		

a) Isolated yield (based upon the capacity of the Fmoc TentaGel S RAM-resin). Determined after correction (by ¹H NMR) for the released PEG in the cleavage step. Purity >95% by GC/MS. b) Irradiated for 6.00 min at 30 W. Purity >80% by GC/MS.

Table 3. Stille and Heck Coupling Reactions on Solid Phase under Microwave Irradiation^a.

a) Isolated yield (based upon the capacity of the Fmoc TentaGel S RAM-resin). Purity >95% by GC/MS. b) The ratio of phenyl vs. butyl transfer in the isolated product was 9:1 according to ¹H NMR. c) 76% conversion according to GC/MS.

COMPARISON OF PRODUCT PATTERN AFTER MICROWAVE AND STANDARD HEATING.

IN SOLUTION: The product pattern after microwave assisted palladium-catalyzed reactions and after standard heating was compared. For the reactions, entry 4 and entry 5 (see TABLE 1 above), were selected for more detailed studies. For the standard heating experiments, there was an optimal initial conversion rate of the arylhalides at 150-170°C. The reaction in entry 4, which represents the most simple system in TABLE 1, is performed with no added ligand and produces a stable product, can be conducted at 150°C in approximately the same reaction times as with microwave technology. A minimum amount of side products are formed with both heating techniques. The reaction in entry 5 represents a more complex system. Standard heating procedures (125-150°C) produces here a very complex product mixture containing less than 20% of the desired product.

ON SOLID SUPPORT: We have compared the product pattern after microwave assisted palladium-catalyzed reactions and after standard heating. Standard heating over 170°C is not compatible with the solid support.

EXPERIMENTAL PART

(*E*)-Methyl-4-methoxycinnamate. Table 1, Entry 1. In the reaction tube were mixed 4-iodoanisole (0.234 g, 1.0 mmol), methyl acrylate (0.108 g, 1.25 mmol), palladium acetate (0.00225 g, 0.01 mmol), tri-*n*-butylamine (0.185 g, 1.0 mmol) and 0.50 ml DMF. under nitrogen. The contents of the flask were irradiated for 3.50 min with an effect of 60 W. After cooling, the product mixture was poured into 25 ml water and was extracted three times with 25 ml DCM. The combined extracts were washed with 25 ml water and concentrated to an oil. The crude product was purified by column chromatography on silica gel using *i*-hexane/diethyl ether (4/1) as the eluent. The yield was 70% (white solid).

(*E*)-Methyl-4-cyanocinnamate. Table 1, Entry 2. In the reaction tube were mixed 4-bromobenzonitrile (0.182 g, 1.0 mmol), methyl acrylate (0.108 g, 1.25 mmol), palladium acetate (0.00225 g, 0.01 mmol), tri-*o*-tolylphosphine (0.0122 g, 0.04 mmol), triethylamine (0.127 g, 1.25 mmol) and 0.50 ml DMF. under nitrogen. The contents of the flask were irradiated for 3.50 min with an effect of 60 W. After cooling, the product mixture was poured into 25 ml water and was extracted three

times with 25 ml DCM. The combined extracts were washed with 25 ml water and concentrated to an oil. The crude product was purified by column chromatography on silica gel using *i*-hexane/diethyl ether (1/1) as the eluent. The yield was 94% (pale yellow crystals).

(*E*)-Stilbene. Table 1, Entry 3. In the reaction tube were mixed iodobenzene (0.204 g, 1.0 mmol), styrene (0.130 g, 1.25 mmol), palladium acetate (0.00225 g, 0.01 mmol), tri-*n*-butylamine (0.185 g, 1.0 mmol) and 0.50 ml DMF under nitrogen. The contents of the flask were irradiated for 2.50 min with an effect of 90 W. After cooling, the product mixture was poured into 25 ml water and was extracted three times with 25 ml DCM. The combined extracts were washed with 25 ml water and concentrated to an oil. The crude product was purified by column chromatography on silica gel using *i*-hexane as the eluent. The yield was 87% (white crystals, terminal/internal arylation 19/1).

1-Naphthylacrylonitrile. Table 1, Entry 4. In the reaction tube were mixed 1-iodonaphthalene (0.254 g, 1.0 mmol), acrylonitrile (0.066 g, 1.25 mmol), palladium acetate (0.00225 g, 0.01 mmol), triethylamine (0.50 ml) and 0.50 ml DMF under nitrogen. The contents of the flask were irradiated for 3.50 min with an effect of 80 W. After cooling, the product mixture was poured into 25 ml water and was extracted three times with 25 ml DCM. The combined extracts were washed with 25 ml water and concentrated to an oil. The crude product was purified by column chromatography on silica gel using *i*-hexane/diethyl ether (2/1) as the eluent. The yield was 90% (white solid, trans/cis 4/1).

2-Phenyl-2,3-dihydrofuran. Table 1, Entry 5. In the reaction tube were mixed iodobenzene (0.102 g, 0.5 mmol), dihydrofuran (0.350 g, 5.0 mmol), palladium acetate (0.00281 g, 0.0125 mmol), potassium acetate (0.147 g, 1.5 mmol), *n*-Bu₄NCI (0.139 g, 0.5 mmol), triphenylphosphine (0.00328 g, 0.0125 mmol) and 1.0 ml DMF under nitrogen. The contents of the flask were irradiated for 6.00 min with an effect of 60 W. After cooling, the product mixture was poured into 25 ml water and was extracted three times with 25 ml DCM. The combined extracts were washed with 25 ml water and concentrated to an oil. The crude product was purified by column chromatography on silica gel using pentane/diethyl ether (19/1) as the eluent. The yield was 58% (yellow oil).

4-*t*-Butylacetophenone. Table 1, Entry 6. In the reaction tube were mixed 4-*t*-butylphenyl triflate (0.282 g, 1.0 mmol), butyl vinyl ether (0.25 g, 2.5 mmol), palladium acetate (0.00561 g, 0.025 mmol), DPPP (0.0206 g, 0.050 mmol) triethylamine (0.121 g, 1.2 mmol) 0.075 ml water and 0.75 ml DMF. under nitrogen. The contents of the flask were irradiated for 2.50 min with an effect of 55 W. After cooling, the product mixture was poured into 25 ml water and was extracted three times with 25 ml diethyl ether. The combined extracts were washed two times with 25 ml brine and concentrated to an oil. The crude product was purified by column chromatography on silica gel using *i*-hexane/diethyl ether (7/1) as the eluent. The yield was 77% (clear oil).

N,N-Dimethyl-2-[2-(2-naphthyl)ethenyloxy]ethanamine. Table 1, Entry 7. In the reaction tube were mixed 2-naphthyl triflate (0.276 g, 1.0 mmol), [2-(dimethylamino)ethoxy]ethene (0.23 g, 2.0 mmol), palladium acetate (0.00673 g, 0.030 mmol), triphenylphosphine (0.0173 g, 0.066 mmol) triethylamine (0.202 g, 2.0 mmol) and 0.75 ml DMF. under nitrogen. The contents of the flask were irradiated for 7.00 min with an effect of 35 W. After cooling, the product mixture was diluted with 50 ml pentane, transferred to a separatory funnel and washed with 2 x 25 ml water. Additional extraction of the aqueous phases was performed with 25 ml pentane. The combined organic portions were then treated 5 times with 20 ml of 0.1 M HCl. The aqueous extracts were combined, poured into a flask containing excess NaOH (1.0 M) and was extracted with 2 x 50 ml pentane. The combined organic phases were dried (K_2CO_3) and concentrated by evaporation. The yield was 87% (yellow oil, trans/cis 1/1).

2-(4-Cyanophenyl)allyltrimethylsilane. Table 1, Entry 8. In the reaction tube were mixed 4-cyanophenyl triflate (0.251 g, 1.0 mmol), allyltrimethylsilane (0.571 g, 5.0 mmol), palladium acetate (0.00673 g, 0.030 mmol), DPPF (0.0366 g, 0.066 mmol), potassium carbonate (0.207 g, 1.5 mmol) and 1.0 ml acetonitrile. under nitrogen. The contents of the flask were irradiated for 5.00 min with an effect of 50 W. After cooling, the product mixture was poured into 25 ml water and was extracted three times with 25 ml diethyl ether. The combined extracts were washed two times with 25 ml brine and concentrated to an oil. The crude product was purified by repeated Kugelrohr distillation. The yield was 54% (clear oil).

2-(2-Naphthyl)*N*-allyldimethylamine. Table 1, Entry 9. In the reaction tube were mixed 2-naphthyl triflate (0.276 g, 1.0 mmol), *N*-allyldimethylamine (0.426 g, 5.0 mmol), palladium acetate (0.00673 g, 0.030 mmol), DPPF (0.0366 g, 0.066 mmol), potassium carbonate (0.207 g, 1.5 mmol) and 1.0 ml acetonitrile under nitrogen. The contents of the flask were irradiated for 5.00 min with an effect of 50 W. After cooling, the product mixture was poured into 25 ml water and was extracted three times with 25 ml diethyl ether. The combined extracts were washed two times with 25 ml brine and concentrated to an oil. The crude product was purified by repeated Kugelrohr distillation. The yield was 64% (clear oil).

4-Methylbiphenyl. Table 1, Entry 10. In the reaction tube were mixed 4-bromotoluene (0.171 g, 1.0 mmol), phenylboronic acid (0.134 g, 1.1 mmol), Pd(PPh₃)₄ (0.0347 g, 0.030 mmol), sodium carbonate (0.212 g, 2.0 mmol), 0.187 ml 95% ethanol, 0.375 ml water and 0.75 ml DME under nitrogen. The contents of the flask were irradiated for 2.50 min with an effect of 55 W. After cooling, the product mixture was poured into 25 ml water and was extracted three times with 25 ml DCM. The combined extracts were washed with 25 ml brine and concentrated to an oil. The crude product was purified by column chromatography on silica gel using *i*-hexane as the eluent. The yield was 55% (white solid).

4'-Phenylacetophenone. Table 1, Entry 11. In the reaction tube were mixed 4-acetylphenyl triflate (0.268 g, 1.0 mmol), phenyltributyltin (0.441 g, 1.2 mmol), Pd₂dba₃ (0.00916 g, 0.010 mmol), triphenylarsine (0.0245 g, 0.080 mmol), lithium chloride (0.127 g, 3.0 mmol) and 1.0 ml NMP under nitrogen. The contents of the flask were irradiated for 2.50 min with an effect of 50 W. After cooling, addition of 1 M aqueous KF (4 ml) with stirring for 120 min, were followed by dilution (DCM) and filtration through a pad of celite. The filtrate was washed with water (25 ml). Additional extraction of the aqueous phase was performed with DCM (2x25 ml). The combined extracts were washed with 25 ml brine and concentrated to an oil. The crude product was purified by column chromatography on silica gel using *i*-hexane/ethyl acetate (10/1) as the eluent. The yield was 68% (white solid).

(*E*)-4-Bromostilbene. Table 1, Entry 12. In the reaction tube were mixed 4-bromoiodobenzene (0.283 g, 1.0 mmol), styrene (0.115 g, 1.1 mmol), palladium acetate (0.00225 g, 0.01 mmol), triethylamine (0.111 g, 1.1 mmol) and 0.50 ml

DMF under nitrogen. The contents of the flask were irradiated for 4.50 min with an effect of 60 W. After cooling, the product mixture was poured into 25 ml water and was extracted three times with 25 ml DCM. The combined extracts were washed with 25 ml water and concentrated to an oil. The crude product was purified by column chromatography on silica gel using *i*-hexane as the eluent. The yield was 63% (white crystals).

General procedure for Suzuki coupling reactions on solid phase. Table 2.

A sealed Pyrex tube was charged, under nitrogen, with 4-iodo or 4-bromo functionalised resin (100 mg, loading ~0.23 mmol/g), $\text{Pd}(\text{PPh}_3)_4$ (1.15 mg, 0.0010 mmol), aryl boronic acid (0.20 mmol), 2M Na_2CO_3 (0.10 ml, 0.20 mmol), H_2O (0.30 ml), ethanol (0.19 ml) and DME (0.75 ml). After irradiation, the mixture was cooled to room temperature. The resin was transferred thereafter to a 3-ml disposable syringe equipped with a porous polyethylene filter using H_2O /DME, washed with successive portions of H_2O , DME, DMF, sat. KCN/DMSO, MeOH, H_2O , MeOH and DCM (2 x 3 ml each) and dried. The resin was treated with 99% aq. TFA for 1 h, filtered and washed with 0.5 ml TFA and 2 x 0.5 ml DCM. The filtrates were combined and evaporated to dryness to yield the biaryl. All products gave satisfactory ^1H NMR spectra as well as appropriate ion identification by mass spectrometry (EI or PDMS) and had purity >95% by GC/MS.

Procedure for the Stille coupling reaction on solid phase. Table 3.


4-Iodo functionalised resin (100 mg, loading ~0.23 mmol/g), Pd_2dba_3 (1.05 mg, 0.00115 mmol), AsPh_3 (1.41 mg, 0.0046 mmol), phenyltributyltin (73.4 mg, 0.20 mmol) and dry NMP (1.0 ml) were placed in a Pyrex tube under nitrogen. The tube was closed, placed in the microwave cavity and irradiated for 5.00 min. at 40 W. After cooling, the product was washed and cleaved from the resin as described for the Suzuki coupling reactions. The ratio of phenyl vs. butyl transfer in the isolated product was 9:1 according to ^1H -NMR (75% yield, white solid).

Procedure for the Heck coupling reaction on solid phase. Table 3.


4-Iodo functionalised resin (50 mg, loading ~0.23 mmol/g), methyl acrylate (0.046 g, 0.53 mmol), palladium acetate (0.00074 g, 0.0033 mmol), tri-*o*-tolylphosphine (0.00322 g, 0.0106 mmol), triethylamine (0.011 g, 0.11 mmol) and 1.0 ml DMF were placed in a Pyrex tube under nitrogen. The tube was closed, placed in the microwave cavity and irradiated for 5.00 min. at 35 W. After cooling, the product was

washed and cleaved from the resin as described for the Suzuki coupling reactions. The crude product was purified by column chromatography on silica gel using *i*-hexane/DCM (1/1) as the eluent. The yield was 71% (white solid).

In the Heck, Suzuki and Stille reactions performed with microwave energy according to the invention catalytic systems commonly used in these reactions are employed. Below a listing of suitable catalytic systems of soluble palladium complexes will follow. The listed catalytic systems are not the active catalysts in the reactions but rather precatalysts generating the active catalyst.



$\text{Pd}(\text{II})\text{OAc}_2$ + ligand (eg. PPh_3 , AsPh_3 , tri(2-furyl)phosphine, tri (*o*-tolyl) phosphine, DPPF (1,1'-bis(diphenylphosphino)ferrocene or BINAP (2,2'-bis(diphenylphosphino)-1,1'-binaphthyl)



Often additions are made to the catalytic systems, such as LiCl , Bu_4NCl , Cu salts, Ag salts, or TI salts.

Furthermore, it is to be understood that the microwave energy used in the method according to the present invention can be generated in different ways. The microwaves should be generated in a controlled and most of all reproducible way. For large volumes (>0.2 mL), this can be achieved by using magnetrons or clystrons. For small reaction volumes (approx. 100 μl) semiconductor generated microwaves are preferably used.

The standing wave, single mode, performance of the applied microwaves makes it possible to focus and optimise the coupling of the microwaves to the reaction samples in a highly reproducible way.

REFERENCES AND NOTES

1. For reviews on the Heck reaction see;
a) Heck, R. F. *Org React.* 1982, 27, 345,
b) Trost, B. M. and Verhoeven, T. R. *Comprehensive Organometallic Chemistry*, Wilkinson, G.; Stone, F. G. A.; Abel, E. W. Eds.; Pergamon Press: Oxford 1982; vol 8, pp. 854, c) Heck, R. F. *Palladium Reagents in Organic Synthesis*; Academic Press; London. 1985; pp 276,
d) Heck, R. F. *Comprehensive Organic Synthesis*; Trost, B. M.; Flemming, I. Eds.; Pergamon Press: Oxford 1991; vol. 4, pp. 833,
e) de Meijere, A.; Meyer, F. E. *Angew. Chem. Int. Ed. Engl.* 1994, 33, 2379,
f) Cabri, W.; Candiani, I. *Acc. Chem. Res.* 1995, 28, 2,
g) Tsuji, J. *Palladium Reagents and Catalysts: Innovations in Organic Chemistry*, John Wiley & Sons, Chichester 1995; pp. 125. For Heck arylation of heteroatom-substituted double bonds see;
h) Daves, G. D. Jr.; Hallberg, A. *Chem. Rev.* 1989, 89, 1433.
 2. For reviews on the Stille reaction see: a) Stille, J. K. *Angew. Chem. Int. Ed. Engl.* 1986, 25, 508, b) Ritter, K. *Synthesis* 1993, 735, c) Tsuji, J. *Palladium Reagents and Catalysts: Innovations in Organic Chemistry*, John Wiley & Sons, Chichester 1995; pp. 228.
 3. For reviews on the Suzuki reaction see:
a) Miyaura, N.; Suzuki, A. *Chem. Rev.* 1995, 95, 2457,
b) Martin, A. R.; Yang, Y. *Acta Chem. Scand.* 1993, 47, 221.
 4. Examples of Suzuki couplings in Combinatorial Chemistry:
a) Frenette, R.; Friesen, R. W. *Tetrahedron Lett.* 1994, 35, 9177,
b) Backes, B. J.; Ellman, J. A. *J. Am. Chem. Soc.* 1994, 116, 11171,
c) Han, Y.; Walker, S. D.; Young, R. N. *Tetrahedron Lett.* 1996, 37, 2703.
- Examples of Heck couplings in Combinatorial Chemistry:
- d) Young, J. K.; Nelson, J. C.; Moore, J. S. *J. Am. Chem. Soc.* 1994, 116,

10841,

- e) Yu, K. L.; Deshpande, M. S.; Vyas, D. *Tetrahedron Lett.* 1994, 35, 8919,
- f) Hiroshige, M.; Hauske, J. R.; Zhou, P. *Tetrahedron Lett.* 1995, 36, 4567,
- g) Goff, D. A.; Zuckermann, R.N. *J. Org. Chem.* 1995, 60, 5748.

Examples of Stille reactions on support-bound aryl halides:

- h) Deshpande, M. S. *Tetrahedron Lett.* 1994, 35, 5613,
- i) Forman, F. W.; Sucholeiki, I. *J. Org. Chem.* 1995, 60, 523.

5. For recent reviews, see:

- a) Hermkens, P. H. H.; Ottenheijm, H. C. J.; Rees, D. *Tetrahedron* 1996, 52, 4527,
- b) Früchtel, J. S.; Jung, G. *Angew. Chem. Int. Ed. Engl.* 1996, 35, 17,
- c) DeWitt, S. H.; Czarnik, A. W. *Acc. Chem. Res.* 1996, 29, 114,
- d) Armstrong, R. W.; Combs, A. P.; Tempest, P. A.; Brown, S. D.; Keating, T. *A. ibid.* 1996, 29, 123,
- e) Ellman, J. A. *ibid.* 1996, 29, 132,
- f) Gordon, E. M.; Gallop, M. A.; Patel, D. V. *ibid.* 1996, 29, 144,
- g) Thompson, L. A.; Ellman, J. A. *Chem. Rev.* 1996, 96, 555, h) Lowe, G. *Chem. Soc. Rev.* 1995, 24, 309.

- 6. Bose, A. K.; Banik, B. K.; Barakat, K. J.; Manhas, M. S.; *Synlett.*, 1993, 575.
- 7. Adamek, F.; Hajek, M. *Tetrahedron Lett.* 1992, 33, 2039.
- 8. Caddick, S. *Tetrahedron* 1995, 51, 10403.
- 9. Heck, R. F.; Nolley, Jr., J. P. *J. Org. Chem.* 1972, 37, 2320.
- 10. Patel, B. A.; Ziegler, C. B.; Cortese, N. A.; Plevyak, J. E.; Zebovitz, T. C.; Terpkko, M.; Heck, R. F. *J. Org. Chem.* 1977, 42, 3903.
- 11. Larock, R. C.; Gong, W. H.; Baker, B. E. *Tetrahedron Lett.* 1989, 30, 2603.
- 12. a) Andersson, C.-M.; Larsson, J.; Hallberg, A. *J. Org. Chem.* 1990, 55, 5757,
b) Larhed, M.; Andersson, C.-M.; Hallberg, A. *Acta. Chem. Scand.* 1993, 47, 212,
c) Larhed, M.; Andersson, C.-M.; Hallberg, A. *Tetrahedron* 1994, 50, 285.

13. a) Miyaura, N.; Yanagi, T.; Suzuki, A. *Synth. Commun.*, 1981, 11, 513,
b) DME was introduced as solvent in the Suzuki reaction by Gronowitz.
Gronowitz, S.; Bobosik, V.; Lawitz, K. *Chem. Scr.* 1984, 23, 120.
14. Farina, V.; Krishnan, B.; Marshall, D. R.; Roth, G. P. *J. Org. Chem.* 1993, 58, 5434.
15. Plevyak, J. E.; Dickerson, J. E.; Heck, R. F. *J. Org. Chem.* 1979, 44, 4078.
16. Melpolder, J. B.; Heck, R. F. *J. Org. Chem.* 1976, 41, 265.
17. a) Cabri, W.; Candiani, I.; Bedeschi, A.; Penco, S.; Santi, R. *J. Org. Chem.* 1992, 57, 1481, b) Cabri, W.; Candiani, I.; Bedeschi, A.; Santi, R. *J. Org. Chem.* 1990, 55, 3654.
18. The 4-halobenzoic acids (4 eq.) were coupled to deprotected resin using 2-(1*H*-benzotriazole-1-yl)-1,1,3,3-tetramethyl-uronium hexafluorophosphate (HBTU, 4 eq.) and DIEA (8 eq.) for 2 h. Remaining amino groups were then capped by acetylation
19. 90 mm Fmoc TentaGel S RAM-resin (S30 023, 0.23 mmol/g capacity) purchased from Rapp Polymere.

CLAIMS

1. A method for transition metal catalyzed organic reactions comprising a heating step performed with microwave energy, **characterized in** that the organic reactions are catalyzed with palladium except Pd/C.

2. A method according to claim 1, **wherein** the soluble palladium catalyzed reactions are:



in which Org is an organic compound and

Org¹ is aryl, heteroaryl, vinyl, acetylenyl, alkyl, allyl, benzyl, acyl, benzoyl, or mono or poly substituted aryl, heteroaryl, vinyl, acetylenyl, alkyl, allyl, benzyl, benzoyl;

X is a halide, triflate, mesitylate, nonaflate, carbonylhalide, sulfonylhalide, perfluoroalkylsulfonate, arylphosphate, alkylphosphate, diarylarsine, diarylphosphine, diarylstibine, arylodonium or diazonium salt;

the double bond in the H-olefin is unsubstituted or mono, di or tri substituted;

R¹ is alkyl, aryl, heteroaryl or mono or poly substituted alkyl, aryl or heteroaryl;

Org² = Org³ and is aryl, heteroaryl, vinyl, acetylenyl, alkyl, allyl, benzyl or mono or poly substituted aryl, heteroaryl, vinyl, acetylenyl, alkyl, allyl, benzyl;

R² is OH, alkoxy, aryloxy or heteroaryloxy.

3. A method according to claims 1 or 2, **wherein** the reactions are performed in solution.

4. A method according to claims 1 or 2, **wherein** the reactions are performed on solid support.
5. A method according to claim 4, **wherein** Org¹, Org² or H-olefin are attached to a solid support.
6. A method according to claim 5, **wherein** Org¹, Org² or H-olefin are attached to a solid support via a linker.
7. A method according to any of the above claims, **wherein** the organic reactions are C-C coupling reactions.
8. A method according to any of the above claims, **wherein** said coupling reactions form part of a combinatorial chemistry process.
9. A method according to any of the above claims, **wherein** the microwave energy is solely or predominantly in the form of a standing wave.
10. Use of microwave-assisted palladium catalyzed reactions for combinatorial chemistry.
11. Use of microwave-assisted palladium catalyzed reactions for creation of chemical libraries.

1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 97/00794

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07B 37/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,N0 classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CA, WPI

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Tetrahedron Letters, Volume 33, No 15, 1992, Frantisek Adámek et al., "Microwave-Assisted Catalytic Addition of Halocompounds to Alkenes" ---	1-11
A	SYNLETT, Volume 8, August 1993, Ajay K. Bose et al., "Simplified Rapid Hydrogenation Under Microwave Irradiation: Selective Transformations of B-Lactams1" page 575 - page 576 --	1-11
A	US 4574038 A (JEFFREY K.S. WAN), 4 March 1986 (04.03.86) -----	1-11

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

19 August 1997

Date of mailing of the international search report

25 -08- 1997

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Eva Johansson
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT
Information on patent family members

06/08/97

International application No.

PCT/SE 97/00794

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
---	---------------------	----------------------------	---------------------

US 4574038 A

04/03/86

NONE

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☒ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.